

BIHEP-TH-97-17 , Nov. 1997

Momentum Spectrum of η' Meson in the Inclusive Decay

$$B \rightarrow \eta' X_s$$

Dong-Sheng Du and Mao-Zhi Yang

CCAST(WORLD LABORATORY) P.O.BOX 8730, BEIJING, 100080

and

Institute of High Energy Physics, Chinese Academy of Sciences,

P.O.Box 918(4), Beijing, 100039, People's Republic of China *

PACS numbers: 13.25.Hw, 13.40.Hq

The momentum distribution for the η' meson produced through the inclusive decay $B \rightarrow \eta' X$ is investigated under two decay mechanisms, $b \rightarrow \eta' s$ and $b \rightarrow \eta' sg$. Although all these two mechanisms can explain the recently observed decay rate of $B \rightarrow \eta' X$, the momentum spectrums for η' meson predicted by them are strongly different. Thus detailed experiment is proposed to distinguish the two cases.

*Mailing address

Recently more attention has been focused on the inclusive and exclusive decay mode of B mesons, $B \rightarrow \eta' X_s$ and $B \rightarrow \eta' K$, which have been observed with relatively large decay rates [1], $Br(B \rightarrow \eta' X_s) = (6.1 \pm 1.6 \pm 1.3) \times 10^{-4}$ and $Br(B \rightarrow \eta' K) = (7.1_{-2.1}^{+2.5} \pm 0.9) \times 10^{-5}$, under the constraint: $2.2 GeV \leq E(\eta') \leq 2.7 GeV$. Many authors have made efforts to explain these conspicuously large decay rates [2–4]. The mechanism via the intermediate formation of a $c\bar{c}$ pair to form the final state containing an η' meson is too weak to explain the observed large decay rate [2]. Besides, there are also mechanisms based on two-body decay of b-quark $b \rightarrow s\eta'$ [2] and three-body decay $b \rightarrow sg\eta'$ [3]. In the two-body decay mechanism, the b-quark decays into s-quark under gluon emission, where the gluon is either on-shell or off-shell, then the gluon transforms into η' meson due to the anomalously large coupling of η' -gluon system [5]. This is a highly non-perturbative process. While in the three-body decay mechanism, the process proceeds via $b \rightarrow sg^*$ with g^* transfers into g and η' because of the anomalously large $\eta' - g^* - g$ coupling [3]. All these two mechanisms can explain the experimental results with suitable coupling constant of η' and gluon field.

In this work we want to study the momentum distribution of η' meson based on the two-body and three-body decay mechanisms $b \rightarrow s\eta'$ and $b \rightarrow sg\eta'$. By comparing the momentum distributions of η' , the two mechanisms can be distinguished in experiment if the momentum spectrum of η' is measured.

For the two-body decay mechanism, Fritzsche has given an ansatz for the effective interaction between the quarks and the gluonic densities [2],

$$\mathcal{H}^{eff} = -const. \frac{G_F}{\sqrt{2}} V_{ts} \frac{\alpha_s}{4\pi^2} \cdot m_b \bar{b}_R s_L (G_{\mu\nu} G^{\mu\nu} + G_{\mu\nu} \tilde{G}^{\mu\nu}), \quad (1)$$

where the const. includes the highly non-perturbative contributions of the strong interactions involved in the decay process. Using the factorization method the amplitudes of $B \rightarrow \eta' X_s$ can be calculated,

$$\langle X\eta' | \mathcal{H}^{eff} | B \rangle = -f_{g*}^{\eta'} \frac{G_F}{\sqrt{2}} V_{ts} \frac{\alpha_s}{4\pi^2} m_b \bar{b}_{RSL}, \quad (2)$$

where $f_{g*}^{\eta'} \equiv \text{const.} \langle \eta' | G_{\mu\nu} G^{\mu\nu} + G_{\mu\nu} \tilde{G}^{\mu\nu} | 0 \rangle$. Here no attempt is made to calculate $f_{g*}^{\eta'}$, because of its strong non-perturbative origin, but to fit the experimental data to get it. With eq.(2), the decay width is obtained,

$$\Gamma_0 = \frac{1}{8\pi} |f_{g*}^{\eta'}|^2 \frac{G_F^2 \alpha_s^2 |V_{ts}|^2}{128\pi^2} |k_c|, \quad (3)$$

where $|k_c| = \frac{1}{2m_b} [(m_b^2 - (m_s + M_{\eta'})^2)(m_b^2 - (m_s - M_{\eta'})^2)]^{1/2}$.

To get the momentum distribution the Fermi motion of the spectator quark inside the B meson should be taken into account. In this work the Fermi motion is considered using the same method in ACCMM model [6]. The spectator quark inside the meson is handled as an on-shell particle, and is attributed with a Fermi motion. Thus the spectator quark has definite mass m_{sp} and its momentum is $|\vec{p}| = p$. The b quark is considered to be off-shell, its virtual mass $M(p)$ can be given in the restframe of the B meson due to the four-momentum conservation as

$$M(p) = \left(M_B^2 + m_{sp}^2 - 2M_B \sqrt{m_{sp}^2 + p^2} \right)^{1/2}. \quad (4)$$

The momentum distribution probability of the Fermi motion $\phi(p)$ is introduced with a Gaussian behaviour in the ACCMM model,

$$\phi(p) = \frac{4}{\sqrt{\pi} p_F^3} e^{-p^2/p_F^2}, \quad (5)$$

where $\frac{4}{\sqrt{\pi}p_F^3}$ is the normalizing constant under the normalization $\int_0^\infty dp p^2 \phi(p) = 1$. The free parameter p_F is taken to be $p_F = 0.57 GeV$, which has been used to explain the momentum spectrum of J/ψ in $B \rightarrow J/\psi X$ [7].

The decay width of b quark should be transformed from the b quark restframe to the B meson restframe with a Lorentz boost,

$$\frac{d\Gamma_b(p_{\eta'}, p)}{dp_{\eta'}} = \begin{cases} 0, & p_{\eta'} < |k_-|, \\ \frac{M_b}{E_b} \frac{\Gamma_0}{k_+ - |k_-|}, & |k_-| < p_{\eta'} < k_+, \\ 0, & p_{\eta'} > k_+ \end{cases} \quad (6)$$

where Γ_0 is expressed in eq.(3) with m_b be substituted with $M(p)$, and $k_\pm = \frac{1}{M_b}(E_b p_0 \pm E_0 p)$, E_0, p_0 are the energy and momentum of η' meson in the rest-frame of b quark, respectively. To compare the momentum distribution with experiment, one has to transform eq.(6) into the laboratory frame where the B meson is in flight. Because the B mesons are produced through the cascade process $e^+e^- \rightarrow \Upsilon(4s) \rightarrow B\bar{B}$, they move with a momentum $p_B = 0.34 GeV$. Performing the Lorentz boost to the laboratory frame, the differential branching ratio for a B meson in flight is,

$$\frac{1}{\Gamma_B} \frac{d\Gamma}{dp_{\eta'}} = \tau_B \int_{|K_-(p_{\eta'})|}^{\hat{K}_+(p_{\eta'})} \frac{dp'_{\eta'}}{k_+(p'_{\eta'}) - |k_-(p'_{\eta'})|} \int_0^{p_{max}} dp p^2 \phi(p) \frac{d\Gamma_b}{dp'_{\eta'}}(p'_{\eta'}, p), \quad (7)$$

where $p_{max} = \frac{1}{2M_B}[(M_B^2 + m_{sp}^2 - M_{\eta'}^2)^2 - 4M_B^2 m_{sp}^2]^{1/2}$, and

$$K_\pm(k) = \frac{1}{M_B}(E_B k \pm p_B E_{\eta'}), \quad \hat{K}_+(k) = \min\{K_+(k), k_{max}\}, \quad (8)$$

k_{max} is the maximum value of η' momentum in the restframe of B meson.

The numerical result is shown by the solid curve of Fig.1. The discussion is given in the end part of the paper.




FIG. 1. Momentum distribution of η' meson in the decay of $B \rightarrow \eta'X$ in the laboratory frame where B meson is in flight. The solid curve is for the $b \rightarrow s\eta'$ case, while the dashed curve for the $b \rightarrow sg\eta'$ case.

Now we turn to the three-body decay mechanism $b \rightarrow sg\eta'$, for the detail of which the readers are referred to Ref. [3]. The loop induced $b \rightarrow s$ current in the standard model is [8]

$$\frac{G_F}{\sqrt{2}} \frac{g_s}{4\pi^2} v_t \bar{s} t^a \{ \Delta F_1 (q^2 \gamma_\mu - q_\mu \not{q}) L - F_2 i \sigma_{\mu\nu} q^\nu m_b R \} b, \quad (9)$$

where $v_t = V_{ts}^* V_{tb}$, $\Delta F_1 = -5.05$, $F_2 \simeq 0.2$. The Feynman rule for the $\eta' - g - g$

vertex is [3],

$$-ia_g c_p \epsilon_{\mu\nu\alpha\beta} \epsilon^\mu(q) \epsilon^\nu(k) q^\alpha k^\beta, \quad (10)$$

here a_g is the effective gluon anomaly coupling. ϵ is the polization of the gluon field, q and k are the four-momentum of the two gluons.

Define $x \equiv \frac{p_x^2}{m_b^2}$, $y \equiv \frac{q^2}{m_b^2}$, $x' \equiv \frac{M_{\eta'}^2}{m_b^2}$, $x'_s \equiv \frac{m_s^2}{m_b^2}$, p_x is the four-momentum of all the inclusive hadronic products except η' meson. The differential branching ratio is

$$\frac{1}{\Gamma_B} \frac{d\Gamma(b \rightarrow sg\eta')}{dp_{\eta'}} = \frac{1}{\Gamma_B} \frac{G_F^2 |v_t|^2 m_b^5}{256\pi^3} \frac{g_s^2}{16\pi^4} \frac{a_g^2}{4} \int dy [\Delta F_1^2 C_1 + F_2 \Delta F_1 C_2 + F_2^2 C_3] \frac{2m_b p_{\eta'}}{E_{\eta'}}, \quad (11)$$

with $C_0 = [-2(x - x'_s)^2 y + (1 - y - x'_s)(y - x')(2x + y - x' - 2x'_s)]/2$, $C_1 = -(1 - y - x'_s)(y - x')^2/y$, and $C_2 = [2(x - x'_s)^2 y^2 - (1 - y - x'_s)(y - x')(2(x - x'_s)y - (1 - x'_s)(y - x'))]/2y^2$, $x = 1 + x' - 2\sqrt{x' + p_{\eta'}^2/m_b^2}$. C_0 , C_1 , C_2 are consistent with these of Ref. [3] (Hou and Tseng) when taking the limit $x'_s \rightarrow 0$.

For the numerical calculation, we take $m_b = 4.8\text{GeV}$, $m_s = 0.175\text{GeV}$, and $a_g = 1.70\text{GeV}^{-1}$ which can ensure that the integrated branching ratio under the constraint $2.2\text{GeV} \leq E_{\eta'} \leq 2.7\text{GeV}$ is the center value of the experimental data. Using eq.(7) to transform eq.(11) into the laboratory frame with B meson in flight, the result of the differential branching ratio is shown by the dashed curve of Fig.1.

Fig.1 shows that in the three-body case (dashed curve), the momentum distribution for η' meson is much broader than that in the two-body decay case (the solid curve). At the same time most part of the curve lies lower than the cut used in the experiment $2.2\text{GeV} \leq E_{\eta'} \leq 2.7\text{GeV}$ ($2.0\text{GeV} \leq p_{\eta'} \leq 2.52\text{GeV}$). While

in the $b \rightarrow \eta' s$ case, no η' meson is found with momentum less than $1.5 GeV$. The two mechanisms can be distinguished by experiment. Thus more detailed experiment is needed.

ACKNOWLEDGEMENTS

This work was supported in part by the China National Nature Science Foundation and the Grant of State Commission of Science and Technology of China.

[1] B. Behrens (CLEO Collaboration), talk presented at the conference on B physics and CP violation, Honolulu, Hawaii, March 1997; F. Wuerthwein (CLEO Collaboration), talk presented at Rencontre de Moriond, QCD and High Energy Hadronic Interactions, Les Arcs, March 1997; D.H. Miller, talk presented at EPS-HEP97, Jerusalem, Israel, Aug. 1997; J. Smith, talk presented at the 1997 Aspen Winter Physics Conference on Particle Physics, January, 1997.

[2] H. Fritzsch, Phys. Lett. B415, 83 (1997).

[3] D. Atwood and A. Soni, Phys. Lett. B405, 150 (1997); W. S. Hou and B. Tseng, Phys. Rev. Lett. 80, 434 (1997); D. Atwood and A. Soni, Phys. Rev. Lett. 79, 5206

(1997).

- [4] I. Halperin and A. Zhitnitsky, Phys. Rev. Lett., 80, 438 (1998); A. Kagan and A. Petrov, hep-ph/9707354.
- [5] H. Fritzsch, M. Gell-Mann and H. Leutwyler, Phys. Lett. B47, 365 (1973); H. Fritzsch and P. Minkowski, Nuovo Cimento A30, 393(1975).
- [6] G. Altarelli, N. Cabibbo, G. Corbò, L. Maiani, and G. Martinelli, Nucl. Phys. B 208, 365 (1982).
- [7] W. F. Palmer, E. A. Paschos, and P. H. Soldan, Phys. Rev. D56, 5794 (1997).
- [8] W. S. Hou, A. Soni and H. Steger, Phys. Rev. Lett. 59, 1521(1987); W.S. Hou, Nucl. Phys. B308, 561 (1988).